



**Scientific Name:** *Novirhabdovirus sp.*

**Discovered by:** Jensen

**Year Named:** 1963

**Common Name:** Viral Hemorrhagic Septicemia (VHS). Also known as 'Egtved virus' in Europe.

**Taxonomy:** Available through ITIS.

**Identification:** VHS is an RNA virus with a bullet-shaped morphology typical of rhabdoviruses and a 11–12 kb nucleotide genome encoding five structural proteins. Viral particles are 170-180 nm in length and 60-70 nm in width (Skall et al. 2005; Elsayad et al. 2006). A classification based on sequences of N- and G-genes reveals four major genotypic groups that correspond to the geographical distribution of the virus: one group includes isolates from European inland waters and northern marine coastal areas, a second group is composed of marine isolates from the Baltic Sea, a third group comprises isolates from the North Sea, and a fourth group comprises North American isolates. Thus far, the virus has been found in Europe, North America, Japan and Korea (Nishizawa et al. 2002; Skall et al. 2005). Based on a comparison of the gene sequences of all North American VHS isolates, Elsayad et al. (2006) propose that the VHS isolate obtained from a muskellunge in Lake St. Clair constitutes a distinct sublineage of the North American genotype that likely originated on the east coast of North America.

The clinical signs of VHS differ depending on the course of infection. In the latent manifestation of the disease, some mortality may occur and fish become hyperactive, sometimes displaying nervous symptoms such as twisting of the body and behavior that involves swimming erratically in circles or in a corkscrew pattern (CFSPH 2003). Conversely, some carriers of the virus may show no symptoms at all (Dopazo et al. 2002). Histopathological changes occur in the liver, kidneys, spleen and skeletal muscle (McAllister 1990); the kidney and spleen appear to be the organs most often targeted by VHS virus (Brudeseth et al. 2002). In the acute form of the disease, fish become lethargic, dark and anemic, with bulging eyes, congested kidneys, mottled liver, and with hemorrhage in the eyes, skin, gills, fin bases, skeletal muscle and viscera (McAllister 1990). Mortality is very high and the disease is short-lived (CFSPH 2003). In the chronic form of the disease, mortality is low and all the symptoms are similar to the acute form, except that hemorrhaging is not common; instead, the liver, spleen and kidneys experience an accumulation of fluid such that the body becomes bloated and the liver and kidneys become very light in color (McAllister 1990). Survivors of infection can be carriers of the virus throughout the rest of their lives.

**Size:** The VHS virus is approximately 170–180 nm long and 60–70 nm wide (Elsayad et al. 2006; McAllister 1990).

**Native Range:** VHS is indigenous to eastern and western Europe, Japan, and the Pacific coast (from California to Alaska) and Atlantic coast of North America. Some evidence suggests that the European strains of VHS are native to the Atlantic Ocean. It is generally believed that all strains of VHS are derived from a common marine ancestor (Skall et al. 2005).

**Nonindigenous Occurrences:** VHS virus has been present in the Great Lakes since at least 2003. The North American strain of the virus was first isolated from muskellunge *Esox masquinongy* caught in the northwest part of Lake St. Clair, Michigan (Elsayad et al. 2006). In 2005, infected freshwater drum *Aplodinotus grunniens* and round goby *Neogobius melanostomus* were captured in the Bay of Quinte, Lake Ontario (Wren and Lee 2006). In the spring and summer of 2006, VHS was detected in fishes in the Thousands Islands area of the St. Lawrence River (Wren and Lee 2006) and in Lake St. Clair, Lake Erie and Lake Ontario (USDA 2006). All isolates from these areas belong to the same sublineage of the North American genotype (M. Faisal, personal communication, November 2006). MI DNR confirmed VHS in Lake Huron in samples of lake whitefish, walleyes and Chinook salmon from 2005.

**Means of Introduction:** It is not known how VHS was initially introduced to the Great Lakes–St. Lawrence River system; however, genetic evidence suggests that the virus originated from the Atlantic coast of North America, possibly via transport in ballast water or infected migratory fishes (Elsayad et al. 2006). Aquaculture activities are implicated in the spread of the virus (Skall et al. 2005; Fisheries Research Services 2006). The potential for transport with bait fish (reviewed by Goodwin et al. 2004) is demonstrated by the virus' recovery in cell culture from frozen Pacific herring *Clupea pallasii* after two freeze/thaw cycles in a conventional freezer (Meyers et al. 1994). Waterfowl might also play a role in transmitting the virus (Peters & Neukirch 1986).

**Status:** The North American strain of VHS virus is present in Lake Huron, Lake St. Clair, Lake Erie, Lake Ontario, and the St. Lawrence River (Elsayad et al. 2006; USDA 2006).

**Ecology:** VHS occurs in both marine and freshwater environments. It requires an incubation period of approximately 7 to 15 days, depending on water temperature (CFSPH 2003). It becomes inactivated in ether, chloroform, glycerol, formalin, sodium hypochlorite, sodium hydroxide, iodophors, UV radiation, or by desiccation, or exposure to pH levels lower than 2.4 or higher than 12.2 (CFSPH 2003; McAllister 1990). VHS is still stable at a pH of 5.0, while the optimum replication pH is 7.4–7.8. The optimum replication temperature is 14–15°C, whereas replication is low at 6°C and almost non-existent at 20°C (De Kinkelin et al. 1980; Bernard et al. 1983; McAllister 1990). The virus becomes inactive after 24 hours at 20°C in water, but can persist for five days at 4°C in water (CFSPH 2003). Consequently, fish mortality from VHS is greatest at 3–12°C and is very rare above 15°C (McAllister 1990).

Fishes are susceptible to infection at any age. VHS is transmitted to juvenile and adult fish most often via urine and sex products that enter a fish through secondary gill lamellae, or possibly through fin bases or via wounds; it cannot enter eggs and infect fish before hatching (McAllister 1990; Brudeseth et al. 2002; CFSPH 2003; Harmache et al. 2006). Juvenile fish are generally more susceptible than adults. Experiments have recorded infection after contact with infected fish and after immersion in infected water; the virus can remain activated in water for several days (McAllister 1990). The VHS virus can persist for long periods of time in the bottom of culture ponds, potentially in invertebrates (CFSPH 2003). There is evidence of infections transferred between farmed and free-living fishes in European inland waters and coastal areas (Stone et al. 1997; Skall et al. 2005). The mortality rate for infected fish varies between 20% and 80%, depending on environmental conditions, and has reached 100% in trout fry (CFSPH 2003).

### **Impact of Introduction**

**A) Realised:** The North American VHS strain is less virulent to salmon and trout than the European strain (Follett et al. 1997) and has not caused large fish kills of these species in the Great Lakes to date (USDA 2006). However, mortality of other species has been documented. In 2005, VHS apparently caused large die-offs of freshwater drum *Aplodinotus grunniens* and round goby *Neogobius melanostomus* in eastern Lake Ontario and muskellunge *Esox masquinongy* in Lake St. Clair (Wren and Lee 2006). In the spring and summer of 2006, VHS was implicated as a cause of large die-offs of round goby and muskellunge in the Thousands Islands area of the St. Lawrence River (Wren and Lee 2006) and die-offs of muskellunge, northern pike *Esox lucius*, gizzard shad *Dorosoma cepedianum*, smallmouth bass *Micropterus dolomieu*, walleye *Sander vitreus* and yellow perch *Perca flavescens* in Lake St. Clair, Lake Erie and Lake Ontario (USDA 2006).

**B) Potential:** Nearly 50 species of fish are known to be susceptible to VHS. The virus was first isolated from most of these species only within the past two decades (Skall et al. 2005). Susceptible fishes include several species of commercial importance (e.g. lake trout *Salvelinus namaycush*, rainbow trout *Oncorhynchus mykiss*, brook trout *Salvelinus fontinalis*, and coregonids *Coregonus* spp.) that have not yet been killed by the virus in the Great Lakes basin. European freshwater-strain VHS infections usually manifest themselves in salmonids, particularly rainbow trout, which suffer high mortality rates. Although the North American strain appears to be of low pathogenicity to salmonids, it has caused mass mortality in a variety of other marine fishes (Kocan et al. 2003; Meyers and Winton 1995; Skall et al. 2005).

### **Remarks:**

A U.S. Federal Order aims to prevent the spread of VHS into aquaculture facilities by restricting the interstate movement and importation of live fish of VHS-susceptible species: [http://www.aphis.usda.gov/vs/aqua/pdf/vhs-fed-order\\_ogc-changes.pdf](http://www.aphis.usda.gov/vs/aqua/pdf/vhs-fed-order_ogc-changes.pdf)

## Voucher Specimens:

## References:

Bernard, J., P. de Kinkelin and M. Bearzotti-Le Berre. 1983. Viral Hemorrhagic Septicemia of rainbow trout: relation between the G polypeptide and antibody production in protection of the fish after infection with the F25 attenuated variant. *Infection and Immunity* 39: 7–14.

Brudeseth, B.E., R.S. Raynard, J.A. King and Ø. Evensen. 2005. Sequential pathology after experimental infection with marine viral hemorrhagic septicemia isolates of low and high virulence in turbot (*Scophthalmus maximus* L). *Veterinary Pathology* 42: 9–18.

Center for Food Security and Public Health (CFSPH). 2003. Viral Hemorrhagic Septicemia. Institute for International Cooperation in Animal Biologics and College of Veterinary Medicine, Iowa State University, Ames, Iowa. 3 pp.

<[http://www.cfsph.iastate.edu/Factsheets/pdfs/viral\\_hemorrhagic\\_septicemia.pdf](http://www.cfsph.iastate.edu/Factsheets/pdfs/viral_hemorrhagic_septicemia.pdf)>

Revision date: 08/08/2005

de Kinkelin, P., M. Bearzotti-Le Berre and J. Bernard. 1980. Viral Hemorrhagic Septicemia of rainbow trout: selection of a thermoresistant virus variant and comparison of polypeptide synthesis with the wild-type virus strain. *Journal of Virology* 36: 652–658.

Dopazo, C.P., I. Bandin, C. López-Vasquez, J. Lamas, M. Noya and J.L. Barja. 2002. Isolation of viral hemorrhagic septicaemia virus from Greenland halibut *Reinhardtius hippoglossoides* caught at the Flemish Cap. *Diseases of Aquatic Organisms* 50: 171–179.

Elsayed, E., M. Faisal, M. Thomas, G. Whelan, W. Batts and J. Winton. 2006. Isolation of viral haemorrhagic septicaemia virus from muskellunge, *Esox masquinongy* (Mitchill), in Lake St. Clair, Michigan, USA reveals a new sublineage of the North American genotype. *Journal of Fish Diseases* 29: 611–619.

Fisheries Research Services. 2006. Risks to wild freshwater fisheries from viral hemorrhagic septicaemia (VHS) disease. Scottish Executive Environment and Rural Affairs Department.

<<http://www.marlab.ac.uk/FRS.Web/Uploads/Documents/VHSVRisksV3.pdf>> Revision date: 31/05/2006

Follett, J.E., T.R. Meyers, T.O. Burton and J.L. Geesin. 1997. Comparative susceptibilities of salmonid species in Alaska to infectious hematopoietic virus (IHNV) and North American viral hemorrhagic septicemia virus (VHSV). *Journal of Aquatic Animal Health* 9: 34–40.

Goodwin, A.E., J.E. Peterson, T.R. Meyers and D.J. Money. 2004. Transmission of exotic fish viruses: the relative risks of wild and cultured bait. *Fisheries* 29(5): 19–23.

- Harmache, A., M. LeBerre, S. Droineau, M. Giovannini and M. Brémont. 2006. Bioluminescence imaging of live infected salmonids reveals that the fin bases are the major portal of entry for *Novirhabdovirus*. *Journal of Virology* 80: 3655–3659.
- Kocan, R., M. Bradley, N. Elder, T. Meyers, W. Batts and J. Winton. 1997. The North American strain of Viral Hemorrhagic Septicemia virus is highly pathogenic for hatchery-reared Pacific herring (*Clupea pallasii*). *Journal of Aquatic Animal Health* 9: 279–290.
- McAllister, P.E. 1990. Fish Disease Leaflet 83. Viral Hemorrhagic Septicemia of Fishes. U.S. Fish and Wildlife Service, National Fisheries Research Center-Leetown, National Fish Health Research Laboratory, Kearneysville, West Virginia.  
<<http://www.lsc.usgs.gov/fhb/leaflets/83.asp>> Revision date: 06/02/2004
- Meyers, T.R., S. Short, K. Lipson, W.N. Batts, J.R. Winron, J. Wilcock and E. Brown. 1994. Association of viral hemorrhagic septicemia virus with epizootic hemorrhages of the skin in Pacific herring (*Clupea harengus pallasii*) from Prince William Sound and Kodiak Island, Alaska, USA. *Diseases of Aquatic Organisms* 19: 27–37.
- Meyers, T.R. and J.R. Winton. 1995. Viral hemorrhagic septicemia virus in North America. *Annual Review of Fish Diseases* 5: 3–24.
- Nishizawa, T., H. Iida, R. Takano, T. Isshiki, K. Nakajima and K. Muroga. 2002. Genetic relatedness among Japanese, American and European isolates of viral hemorrhagic septicemia virus (VHSV) based on partial G and P genes. *Diseases of Aquatic Organisms* 48: 143–148.
- Peters, F. and M. Neukirch. 1986. Transmission of some fish pathogenic viruses by the heron, *Ardea cinerea*. *Journal of Fish Diseases* 9: 539–544.
- Skall, H.F., N.J. Olesen and S. Møllergaard. 2005. Viral Hemorrhagic Septicaemia Virus in marine fish and its implications for fish farming – a review. *Journal of Fish Diseases* 28: 509–529.
- Stone, D.M., K. Way and P.F. Dixon. 1997. Nucleotide sequence of the glycoprotein gene of Viral Hemorrhagic Septicemia (VHS) viruses from different geographical areas: a link between VHS in farmed fish species and viruses isolated from North Sea cod (*Gadus morhua* L.). *Journal of General Virology* 78: 1319–132.
- United States Department of Agriculture (USDA). 2006. Viral Hemorrhagic Septicemia in the Great Lakes: July 2006 Emerging Disease Notice. Animal and Plant Health Inspection Service. <[http://www.aphis.usda.gov/vs/ceah/cei/taf/emergingdiseasenotice\\_files/vhsgreatlakes.htm](http://www.aphis.usda.gov/vs/ceah/cei/taf/emergingdiseasenotice_files/vhsgreatlakes.htm)> Revision date: 8/12/2006

Wren, M. and S. Lee. 2006. DEC Confirms virus in Lake Ontario and St. Lawrence River fish; Cornell University, USGS document cases of Viral Hemorrhagic Septicemia. New York State NEWS, Department of Environmental Conservation, New York.  
<<http://www.dec.state.ny.us/website/press/pressrel/2006/200683.html>> Revision date: 13/06/2006

---

**Authors:** Rebekah M. Kipp and Anthony Ricciardi

**Revision Date:** December 8, 2006.

**Citation for this Information:** Rebekah M. Kipp and Anthony Ricciardi 2006. GLANSIS.

**Group:** Does not fit groups available

**Lake(s):** Lake Ontario, Lake Erie, Lake St. Clair

**Genus:** *Novirhabdovirus*

**Species:** Undescribed

**Common Name:** Viral Hemorrhagic Septicemia (VHS), Egtved disease.

**Status:** Reported

**Freshwater/Marine:** All

**Pathway:** Unknown

**Exotic/Transplant:** Native Transplant